

Efficacy of Albendazole and Mebendazole With or Without Levamisole for Ascariasis and Trichuriasis

by Turnitin ®

Submission date: 16-Dec-2023 08:02AM (UTC+0700)

Submission ID: 2260354569

File name: OAMJMS-7-1299.pdf (197.33K)

Word count: 3469

Character count: 20067

4 Efficacy of Albendazole and Mebendazole With or Without Levamisole for Ascariasis and Trichuriasis

Endy Juli Anto¹, Sony Eka Nugraha^{2*}

¹Department of Parasitology, Faculty of Medicine, University of Methodist Indonesia, Medan, Indonesia; ²Department of Pharmaceutical Biology, Faculty of Pharmacy, Universitas Sumatera Utara, Medan, Indonesia

Citation: Anto EJ, Nugraha SE. Efficacy of Albendazole and Mebendazole With or Without Levamisole for Ascariasis and Trichuriasis. Open Access Maced J Med Sci. 2019 Apr 30; 7(8):1299-1302. <https://doi.org/10.3889/oamjms.2019.299>

Keywords: Soil-transmitted helminths; *T. trichiura*; Albendazole; Albendazole-levamisole; Mebendazole-levamisole

***Correspondence:** Sony Eka Nugraha, Department of Pharmaceutical Biology, Faculty of Pharmacy, Universitas Sumatera Utara, Medan, Indonesia. E-mail: sonyekanugraha@usu.ac.id

Received: 10-Feb-2019; **Revised:** 17-Apr-2019; **Accepted:** 18-Apr-2019; **Online first:** 28-Apr-2019

Copyright: © 2019 Endy Juli Anto, Sony Eka Nugraha. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

Abstract

BACKGROUND: Helminthiasis in school-aged children potentially causing physical growth and intellectual development retardation. Trichuriasis was the most common type of helminthiasis in children.

AIM: To investigate the efficacy and side effects of albendazole, albendazole combined with levamisole and mebendazole combined with levamisole for trichuriasis and ascariasis.

METHODS: This study was conducted as a double-blind, randomised clinical trial by comparing the efficacy and side effects of albendazole, albendazole combined with levamisole and mebendazole combined with levamisole for trichuriasis. The sample of this study were 180 elementary school students at Deli Serdang Regency State Elementary School, Medan, Indonesia. The study was conducted from April to June 2015.

RESULT: The cure rate of helminthiasis on the 7th day was 81.7% after albendazole therapy, 88.3% after albendazole levamisole therapy, and 83.3% after mebendazole combined with levamisole therapy ($p = 0.577$). Cure rate on the 14th day was 88.3%, 95%, and 91.7% for albendazole, albendazole combined with levamisole, and mebendazole combined with levamisole therapy, respectively ($p = 0.418$). On the 21th day, the cure rate was 88.3%, 96.7%, and 91.7% ($p = 0.230$). Combination of albendazole and levamisole showed the highest cure rate, despite the statistically insignificant difference for all groups ($p > 0.05$). Combination of albendazole combined with levamisole showed better cure rate for mild trichuriasis (95.8%) than albendazole therapy (46.2%) and mebendazole combined with levamisole (83.3%), ($p < 0.05$).

CONCLUSION: Single-dose albendazole, a combination of albendazole and levamisole, and a combination of mebendazole and levamisole had similar efficacy in reducing egg count in helminthiasis. Combination of albendazole and levamisole showed better cure rate for mild trichuriasis and mixed infections. Side effects were similar in all treatment groups.

Introduction

Intestinal worm infection is still a major public health problem in Indonesia, especially in rural areas. In Indonesia and other developing countries, *Ascaris lumbricoides*, *Trichuris trichiura*, and hookworm are the most common intestinal parasites [1], [2]. World Health Organization (WHO) estimated that at least two billion people or nearly one-third of the world's population had been infected with soil-transmitted helminths (STH) or helminthiasis [3]. About 300 million infected people suffered from severe illness, and about 400 million school-aged children worldwide had

been infected [4], [5].

Many species of worms had been reported to cause infection in Indonesia. *Trichuris trichiura*, was resided in the human caecum, was the most common cause of helminthiasis and one of the most important intestinal nematodes in human [6], [7]. School-age children were frequently infected with helminthiasis, which potentially causing diarrhoea, nutritional deficiency, anaemia, growth disorders, and intellectual disturbance [8], [9].

Public health program to control helminthiasis was largely dependent on the administration of anti-helminthic drugs for elementary-school children [10].

Theoretically, there were several broad-spectrum anti-helminthic drugs such as albendazole, levamisole, mebendazole, and pyrantel pamoate with variable advantages and disadvantages [11], [12], [13], [14]. Benzimidazole group was the most common anti-helminthic used in a public health setting [15], [16], [17].

We conducted a clinical trial to compare the efficacy and side effects of albendazole, albendazole with levamisole and mebendazole with levamisole therapy against trichuriasis and ascariasis in elementary-school children to determine the most effective regimen.

Material and Methods

This study was a double-blind, randomised clinical trial comparing the efficacy and side effects of albendazole, albendazole combined with levamisole and mebendazole combined with levamisole for trichuriasis. The sample of this study were 180 elementary-school students at Deli Serdang Regency State Elementary School, Medan, Indonesia. The study was conducted from April to June 2015. Study steps consisted of preparation, data collection, processing, data analysis, and data improvement.

Written informed consent was conducted by all subjects' parents or guardian two weeks before the scheduled intervention date. A container was given to each subject to keep their stool sample at home. The collected stool samples were qualitatively and quantitatively examined by the Kato-Katz method at Parasitology Laboratory of Medical Faculty, University of Sumatera Utara, Medan. Stool sample with a positive finding of trichuriasis was randomised as the study subjects into the three treatment groups. Stool sample examination for evaluation was done on day 7, day 14 and 21 after treatment.

Each subject in the first group was given one tablet of 400 mg albendazole. Second group was given one tablet of 400 mg albendazole and 2 tablets of 25 mg levamisole (for children weighed 10 – 20 kg), or one tablet of 400 mg albendazole and 4 tablets of 25 mg levamisole (for children weighed 21 – 40 kg), or one tablet of 400 mg albendazole and 6 tablets of 25 mg levamisole (for children weighed > 40 kg). The third group was given one tablet of 500 mg mebendazole and levamisole dose is the same as the second group. For the following 14 days, parents were instructed to record any side effect after treatment. Comparison of recovery rate between the intervention groups was analysed using chi-square test or Wilcoxon test as suitable. The p-value of less than 0.05 was considered significant.

Results

There were 807 Deli Serdang elementary students; after the exclusion, we found 185 students with helminthiasis based on stool examination. Five of them had started therapy before the study. Therefore, they were excluded. The remaining 180 students were study subjects who undergone randomisation into three intervention groups (60 students in each group). The characteristics of the subjects can be seen in Table 1.

Table 1: Characteristics of Subjects

Characteristics	First Group Albendazole (N = 60)	Second Group Albendazole + Levamisole (N = 60)	Third Group Mebendazole + Levamisole (N = 60)
Age, years (mean \pm SD)	9.2 \pm 1.734	8.9 \pm 1.540	9.1 \pm 1.567
Sex, n (%)			
Male, n (%)	35 (58.3)	28 (46.7)	31 (51.7)
Female, n (%)	25 (41.7)	32 (53.3)	29 (48.3)
Weight, kg	27.2 \pm 7.957	27.1 \pm 6.374	27.9 \pm 7.363
Height, cm (mean \pm SD)	129.5 \pm 13.944	132.0 \pm 18.516	128.4 \pm 12.785
Weight/height (mean \pm SD)	97.1 \pm 6.574	95.7 \pm 13.014	94.1 \pm 17.757
Parental occupation			
Entrepreneur	38 (16.7)	14 (23.3)	16 (26.6)
Farmer	11 (18.3)	45 (75)	38 (63.3)
Civil servant	1 (1.6)	-	1 (1.6)
Others	10 (16.6)	1 (1.6)	5 (8.3)
Parental education, n (%)			
Uneducated	1 (0.83)	7 (5.83)	13 (10.83)
Elementary school	22 (18.33)	36 (30)	25 (20.83)
Junior high school	46 (38.33)	38 (31.67)	42 (35)
Senior high school	3 (35.83)	39 (32.5)	38 (31.67)
University	8 (6.67)	-	2 (1.67)
Helminthiasis (%)			
A. lumbricoides	47 (78.3)	34 (56.7)	11 (18.3)
T. trichura	6 (10)	19 (31.6)	12 (20)
A. lumbricoides + T. trichura	7 (11.7)	7 (11.7)	36 (60)
A. lumbricoides + T. trichura + Enterobius	-	-	1 (1.7)

The most common helminthiasis was infection by *Ascaris lumbricoides*, followed by mixed infection by *Ascaris lumbricoides* with *Trichuris trichura*, *Trichuris trichura* only, and mixed infection by *Ascaris lumbricoides* with *Trichuris trichura* and *Enterobius vermicularis* as seen in Table 2.

Table 2: Prevalence of helminthiasis

Helminthiasis Etiology	Numbers (%)
<i>Ascaris lumbricoides</i>	92 (51.11)
<i>Trichuris trichura</i>	37 (20.55)
A. lumbricoides + T. trichura	50 (27.76)
A. lumbricoides + T. trichura + E. vermicularis	1 (0.55)

Subjects with mild *A. lumbricoides* infection were 50 children (83.3%) in the first group, 39 students (65%) in the second group, and 46 students (76.7%) in the third group. Subjects with mild *T. trichura* infection were 13 children (21.7%) in the first group, 24 children (40%) in the second group, and 24 children (40%) in the third group.

Table 3: Basic Characteristics of Research Based on Intensity of Infection

Characteristics	First Group Albendazole (N = 60)	Second Group Albendazole + Levamisole (N = 60)	Third Group Mebendazole + Levamisole (N = 60)
The intensity of infection, n (%)			
<i>Ascaris lumbricoides</i>			
Mild	50 (83.3)	39 (65)	46 (76.7)
Moderate	4 (6.7)	2 (65)	1 (1.7)
<i>Trichuris trichura</i>			
Mild	13 (21.7)	24 (40)	24 (40)
Moderate	-	2 (3.3)	1 (1.7)

The Statistical analysis did not show a significant reduction in egg count on the 7th day after therapy in all groups, whereas there was a significant reduction in the 14th and 21st day in all groups. Determination of egg reduction rate can be seen in Table 4.

Table 4: Egg Reduction Rate on Day 7th, 14th, 21th

Parasites	Anthelmintics regimen	Total eggs per gram			P value
		Mean (SD) 7 th day	Mean (SD) 14 th day	Mean (SD) 21 st day	
A. lumbricoides	Albendazole	34.00 ± 165.64	-	-	0.651
	Albendazole + Levamisole	14.40 ± 111.54	-	-	
	Mebendazole + Levamisole	34.80 ± 126.25	-	-	
	Mebendazole + Levamisole	34.80 ± 126.25	-	-	
T. trichiura	Albendazole	33.60 ± 95.91	24.80 ± 81.23	20.40 ± 74.96	0.247
	Albendazole + Levamisole	20.00 ± 89.38	12.40 ± 73.47	7.60 ± 47.89	
	Mebendazole + Levamisole	40.00 ± 122.66	18.80 ± 68.03	8.80 ± 29.63	
	Mebendazole + Levamisole	40.00 ± 122.66	18.80 ± 68.03	8.80 ± 29.63	

The cure rate of helminthiasis on the 7th day was 81.7% after albendazole therapy, 88.3% after albendazole + levamisole therapy, and 83.3% after mebendazole + levamisole therapy (p = 0.577). Cure rate on the 14th day was 88.3%, 95%, and 91.7% for albendazole, albendazole + levamisole, and mebendazole + levamisole therapy, respectively (p = 0.418). On the 21st day, the cure rate was 88.3%, 96.7%, and 91.7% (p = 0.230). Combination of albendazole and levamisole showed the highest cure rate, despite the statistically insignificant difference for 8 groups (p > 0.05). Determination of cure rates analysis can be seen in Table 5.

Table 5: The Cure Rates Analysis On Day 7th, 14th, 21th

Recovery Therapy	Recovered		Not Recovered		P
	n	%	n	%	
Albendazole (Day-7)	49	81.7	11	18.3	0.577
Albendazole + Levamisole (Day-7)	53	88.3	7	11.7	
Mebendazole + Levamisole (Day-7)	50	83.3	10	16.7	
Albendazole (Day-14)	53	88.3	7	11.7	0.418
Albendazole + Levamisole (Day-14)	57	95.0	3	5.0	
Mebendazole + Levamisole (Day-14)	55	91.7	5	8.3	
Albendazole (Day-21)	53	88.3	7	11.7	0.230
Albendazole + Levamisole (Day-21)	58	96.7	2	3.3	
Mebendazole + Levamisole (Day-21)	55	91.7	5	8.3	

Combination of albendazole combined with levamisole showed better cure rate for mild trichuriasis (95.8%) than albendazole therapy (46.2%) and mebendazole + levamisole (83.3%), p 0.00. Determination of cure rate mild helminthiasis analysis can be seen in Table 6 and Table 7.

Table 6: The Cure Rate of Mild Helminthiasis

Parasite(s)	Treatment	Cured		Recovery		P
		n	%	n	%	
A. lumbricoides	Albendazole	46	92	4	8	0.176
	Albendazole + Levamisole	39	100	-	-	
	Mebendazole + Levamisole	42	91.3	4	8.7	
T. trichiura	Albendazole	6	46.2	7	53.8	0.01
	Albendazole + Levamisole	23	95.8	1	4.2	
	Mebendazole + Levamisole	20	83.3	4	16.7	

The side effects during the treatment process in each group had been observed. Side effect observed in the albendazole group were 13.3%, Albendazole combined with Levamisole was 26.7%,

and Mebendazole combined with Levamisole were 20 %.

Table 7: Cure Rate of Each Intervention Group

Parasite(s)	Intervention	Cure Rate (%)	P Value
A. lumbricoides	Albendazole	100	-
	Albendazole + Levamisole	100	
	Mebendazole + Levamisole	100	
T. trichiura	Albendazole	66.7	0.136
	Albendazole + Levamisole	94.7	
	Mebendazole + Levamisole	92.3	
A. lumbricoides + T. trichiura	Albendazole	28.6	0.079
	Albendazole + Levamisole	85.7	
	Mebendazole + Levamisole	66.7	

Observation data of side effect can be seen in

Table 8

Table 8: Side Effects analysis

Side Effect	Albendazole n (%)	Albendazole + Levamisole n (%)	Mebendazole + Levamisole n (%)
None	52 (86.7)	44 (73.3)	48 (80.0)
Yes	8 (13.3)	16 (26.7)	12 (20.0)

Discussion

Helminthiasis has still been a major health problem in Indonesia. A. lumbricoides, T. trichiura and hookworm (N. americanus and A. duodenale) were the most common aetiology. WHO data on 2012 reported a high prevalence of helminthiasis in North Sumatera, i.e. 80% of school-aged children [18], [19].

Statistical analysis did not show a significant reduction in egg count on the 7th day after therapy in all groups, whereas there was a significant reduction in the 14th and 21st day in all groups. Even after the reduction in egg number at 14th and 21st day, we still found several T. trichiura eggs in subjects' stool, indicating the difficulty in eradicating trichuriasis as mentioned in the literature [20]. A study by Saputri in 2010 found significant egg reduction rate in single-dose mebendazole and mebendazole with levamisole therapy for A. lumbricoides and T. trichiura infections [21]. The contradictive result was found by Sihite et al., (2014) and Knopp et al., (2010) study, which found no significant difference in the treatment with mebendazole, albendazole, and mebendazole with levamisole [22], [23].

Based on Table 5, the combination of albendazole and levamisole showed the highest cure rate, despite the statistically insignificant difference for all groups (p > 0.05). Therefore, this finding indicated the similar efficacy of albendazole, albendazole + levamisole, and mebendazole + levamisole therapy.

Based on Table 6, the combination of albendazole with levamisole showed better cure rate for mild trichuriasis (95.8%) than albendazole therapy (46.2%) and mebendazole combined with levamisole (83.3%), p = 0.01. We hypothesised that levamisole had enhanced efficacy than albendazole and mebendazole for mild trichuriasis. Previous studies

showed that a single dose of albendazole or mebendazole had 28% and 36% recovery rate, respectively [16], [23]. For mixed infection, albendazole combined with levamisole was more effective (cure rate 85.7%) than single albendazole (28.65%) or mebendazole combined with levamisole (66.7%), $p = 0.079$ that showed no significant difference in the recovery of helminthiasis with mebendazole + levamisole or single mebendazole therapy [22]. Another study found no significant differences in the recovery rate of helminthiasis between mebendazole with or without levamisole therapy [21]. The most common side effects in all groups were nausea and diarrhoea. No serious side effects were observed in this study, and mild side effects had recovered by their own. Table 8 shows that there was no difference in side effects between intervention groups.

It can be concluded that single-dose albendazole, a combination of albendazole and levamisole, and the combination of mebendazole and levamisole had similar efficacy in reducing egg count in helminthiasis. Combination of albendazole and levamisole showed better cure rate for mild trichuriasis and mixed infections. Side effects were similar in all treatment groups.

References

1. Sakti H, Nokes C, Hertanto W, Hendratno S, Hall A, Bundy DAP, et al. Evidence for an association between hookworm infection and cognitive function in Indonesian school children. *Tropical Medicine and International Health*. 1999; 4(5):322-34. <https://doi.org/10.1046/j.1365-3156.1999.00410.x> PMID:10402967
2. Toma T, Miyagi I, Hasegawa H, Kamimura K, Tokuyama Y, Selomo M, et al. Prevalence of intestinal helminth infections in Barru, S, Sulawesi, Indonesia. *Parasitology International*. 1998; 47:323. [https://doi.org/10.1016/S1383-5769\(98\)80943-4](https://doi.org/10.1016/S1383-5769(98)80943-4)
3. World Health Organization. Soil-transmitted helminthiasis: number of children treated in 2014. *Weekly Epidemiological Record*. 2015; 90(51/52):705-11.
4. World Health Organization. Schistosomiasis and soiltransmitted helminth infections preliminary estimates of the number of children treated with albendazole or mebendazole. *Weekly Epidemiological Record*. 2006; 81(16):145-63.
5. Steinmann P, Utzinger J, Du ZW, Jiang JY, Chen JX, Hattendorf J, Zhou H, Zhou XN. Efficacy of single-dose and triple-dose albendazole and mebendazole against soil-transmitted helminths and *Taenia* spp.: a randomized controlled trial. *PLoS one*. 2011; 6(9):e25003. <https://doi.org/10.1371/journal.pone.0025003>
6. De Silva NR, Brooker S, Hotez PJ, Montresor A, Engels D, Savioli L. Soil-transmitted helminth infections: updating the global picture. *Trends in Parasitology*. 2003; 19(12):547-51. <https://doi.org/10.1016/j.pt.2003.10.002> PMID:14642761
7. Drake LJ, Jukes MC, Sternberg RJ, Bundy DA. Geohelminth infections (ascariasis, trichuriasis, and hookworm): cognitive and developmental impacts. *Seminars in Pediatric Infectious Diseases*. 2000; 11(4):245-251. <https://doi.org/10.1053/spid.2000.9638>
8. Albonico M, Allen H, Chitsulo L, Engels D, Gabrielli AF, Savioli L. Controlling soil-transmitted helminthiasis in pre-school-age children through preventive chemotherapy. *PLoS neglected tropical diseases*. 2008; 2(3):e126. <https://doi.org/10.1371/journal.pntd.0000126>
9. Amelia F, Ali M, Pasaribu S. Mebendazole vs. mebendazole-pyranterl pamoate for soil-transmitted helminthiasis infection in children. *Paediatrica Indonesiana*. 2013; 53(4):209-13. <https://doi.org/10.14238/pi53.4.2013.209-13>
10. World Health Organization. Report of the WHO Informal Consultation on the Use of Chemotherapy for the Control of Morbidity Due to Soil-Transmitted Nematodes in Humans, Geneva, 29 April to 1 May 1996. Geneva: World Health Organization, 1996.
11. Chavarria Ap, Swartzwelder Jc, Zeledón R, Villarejos Vm. Mebendazole, An Effective Broad-Spectrum Anthelmintic. *The American Journal of Tropical Medicine and Hygiene*. 1973; 22(5):592-5. <https://doi.org/10.4269/ajtmh.1973.22.592> PMID:4729739
12. Krepel HP, Haring T, Baeta S, Polderman AM. Treatment of mixed Oesophagostomum and hookworm infection: effect of albendazole, pyrantel pamoate, levamisole and thiabendazole. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1993; 87(1):87-9. [https://doi.org/10.1016/0035-9203\(93\)90437-U](https://doi.org/10.1016/0035-9203(93)90437-U)
13. Ramalingam S, Sinniah B, Krishnan U. Albendazole, an effective single dose, broad spectrum anthelmintic drug. *The American journal of tropical medicine and hygiene*. 1983; 32(5):984-9. <https://doi.org/10.4269/ajtmh.1983.32.984> PMID:6625078
14. Olliaro P, Seiler J, Kuesel A, Horton J, Clark JN, Don R, Keiser J. Potential drug development candidates for human soil-transmitted helminthiasis. *PLoS neglected tropical diseases*. 2011; 5(6):e1138. <https://doi.org/10.1371/journal.pntd.0001138>
15. Martin RJ, Robertson AP, Bjorn H. Target sites of anthelmintics. *Parasitology*. 1997; 114(7):111-24.
16. Keiser J, Utzinger J. Efficacy of current drugs against soil-transmitted helminth infections: systematic review and meta-analysis. *Jama*. 2008; 299(16):1937-48. <https://doi.org/10.1001/jama.299.16.1937> PMID:18430913
17. Cook GC. Use of benzimidazole chemotherapy in human helminthiasis: indications and efficacy. *Parasitology Today*. 1990; 6(4):133-6. [https://doi.org/10.1016/0169-4758\(90\)90232-S](https://doi.org/10.1016/0169-4758(90)90232-S)
18. World Health Organization. Research priorities for helminth infections. WHO Technical Report Series. World of Health Organization Bulletin; 14, 2012.
19. Lubis IN, Pasaribu S, Lubis CP. Current status of the efficacy and effectiveness of albendazole and mebendazole for the treatment of *Ascaris lumbricoides* in North-Western Indonesia. *Asian Pacific journal of tropical medicine*. 2012; 5(8):605-9. [https://doi.org/10.1016/S1995-7645\(12\)60125-4](https://doi.org/10.1016/S1995-7645(12)60125-4)
20. Hall A, Nahar Q. Albendazole and infections with *Ascaris lumbricoides* and *Trichuris trichiura* in children in Bangladesh. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1994; 88(1):110-2. [https://doi.org/10.1016/0035-9203\(94\)90525-8](https://doi.org/10.1016/0035-9203(94)90525-8)
21. Dewi S. Comparison of the Efficacy of a Single Dose of Mebendazole with and without Levamisole on Transmitted Soil Worms in Elementary School-aged Children. Thesis: Medan, 2009. Available online: <http://repository.usu.ac.id/handle/123456789/16639> (accessed on 8 December 2016)
22. Sihite IF, Ali M, Pasaribu AP, Pasaribu S, Lubis CP. Efficacy of mebendazole and levamisole, alone or in combination, for soil-transmitted helminthiasis. *Paediatrica Indonesiana*. 2014; 54(1):9-14. <https://doi.org/10.14238/pi54.1.2014.9-14>
23. Knopp S, Mohammed KA, Speich B, Hattendorf J, Khamis IS, Khamis AN, Stothard JR, Rollinson D, Marti H, Utzinger J. Albendazole and mebendazole administered alone or in combination with ivermectin against *Trichuris trichiura*: a randomized controlled trial. *Clinical infectious diseases*. 2010; 51(12):1420-8. <https://doi.org/10.1086/657310> PMID:21062129

Efficacy of Albendazole and Mebendazole With or Without Levamisole for Ascariasis and Trichuriasis

ORIGINALITY REPORT

11%

SIMILARITY INDEX

9%

INTERNET SOURCES

9%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to TAFE Queensland Brisbane Student Paper	2%
2	www.neliti.com Internet Source	1%
3	www.science.gov Internet Source	1%
4	academicctree.org Internet Source	1%
5	"Abstracts—APASL 2013", Hepatology International, 2013 Publication	1%
6	Yulia Fatma Wardani, Ida Safitri Laksono, Teti Adriana Lubis. "Trichuris dysentery syndrome, the neglected tropical disease: a case series", Paediatrica Indonesiana, 2022 Publication	1%
7	Adrian, Rony Abdi Syahputra, Sukirman Lie, Steven Theo, Sony Eka Nugraha. "Antioxidant, Total Phenol, Total Flavonoid, and LC-MS/MS	1%

Analysis of Pometia Pinnata Ethanol Extract",
2021 IEEE International Conference on
Health, Instrumentation & Measurement, and
Natural Sciences (InHeNce), 2021

Publication

8	journal.uniku.ac.id Internet Source	1 %
9	espace.library.uq.edu.au Internet Source	1 %
10	DANIEL KULKE, JÜRGEN KRÜCKEN, ACHIM HARDER, RALPH KREBBER, KRISTINE FRAATZ, HEINZ MEHLHORN, GEORG VON SAMSON-HIMMELSTJERNA. " efficacy of PF1022A and nicotinic acetylcholine receptor agonists alone and in combination against ", Parasitology, 2013 Publication	1 %
11	Lloyd, A. E., B. L. Honey, B. M. John, and M. Condren. "Treatment Options and Considerations for Intestinal Helminthic Infections", Journal of Pharmacy Technology, 2014. Publication	1 %
12	Submitted to Universitas Sumatera Utara Student Paper	1 %
13	ojbe.org Internet Source	1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On