

ORIGINALITY REPORT

20%

SIMILARITY INDEX

16%

INTERNET SOURCES

8%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

www.hindawi.com

Internet Source

3%

2

worldwidescience.org

Internet Source

2%

3

caspjim.com

Internet Source

2%

4

www.wjgnet.com

Internet Source

2%

5

www.e-sciencecentral.org

Internet Source

2%

6

Uygur-Bayramiçli, Oya. "A clinical dilemma: abdominal tuberculosis", World Journal of Gastroenterology, 2003.

Publication

2%

7

D. Santoso, E.J. Wahyudi, W.G.A. Kadir, S. Alawiyah, A. D. Nugraha, P. Supendi, W.W. Parnadi. "Gravity Structure around Mt. Pandan, Madiun, East Java, Indonesia and Its Relationship to 2016 Seismic Activity", Open

1%

8	www.indianjmedsci.org Internet Source	1%
9	Emre Ozgu. "Tuberculosis in Gynecologic Oncology: Still a Diagnostic Dilemma", Obstetrics & Gynecology International Journal, 2014 Publication	1%
10	Boujaoude, Joseph D. "Diagnosis by endoscopic ultrasound guided fine needle aspiration of tuberculous lymphadenitis involving the peripancreatic lymph nodes: A case report", World Journal of Gastroenterology, 2007. Publication	1%
11	icime.usu.ac.id Internet Source	1%
12	documents.mx Internet Source	1%
13	de.scribd.com Internet Source	1%

Prolonged_Fever_in_Peritoneal_TB-compressed.pdf

by

Submission date: 23-Dec-2020 07:49AM (UTC-0500)

Submission ID: 1480853007

File name: Prolonged_Fever_in_Peritoneal_TB-compressed.pdf (110.77K)

Word count: 1548

Character count: 9198

PAPER • OPEN ACCESS

Prolonged fever in peritoneal tuberculosis: A case report

To cite this article: U Zein *et al* 2018 *IOP Conf. Ser.: Earth Environ. Sci.* **125** 012081

View the [article online](#) for updates and enhancements.

Related content

- 2** - [Prolonged fever and involuntary weight loss as manifestations of bacterial endocarditis: A case report](#)
N M D D Sukmawati, T P Merati, A Somia et.al.
- 2** - [Case report : tuberculosis liver abscess in male alcoholism patient](#)
W P Siahaan and F Ginting
- 11** - [Late diagnosis of multiple myeloma: a case report](#)
A Syahreza, D Gatot and A I Mardia

Prolonged fever in peritoneal tuberculosis: A case report

U Zein^{1,2*}, S Irwandi², H Habib¹, H Lim³, M Pasha¹, I Janis⁴, R H Saragih⁵, Y Ginting⁵ and R Effendy-Y S⁶

¹Dr.Umar Zein Tropical Disease and Infectious Clinic, Medan, Indonesia

²Department of Internal Medicine, Faculty of Medicine, Universitas Islam Sumatera Utara, Medan, Indonesia

³Department of Pharmacology, Faculty of Medicine, Universitas Methodist, Medan, Indonesia

⁴Department of Microbiology, Faculty of Medicine, Universitas Islam Sumatera Utara, Medan, Indonesia

⁵Division of Tropical Diseases and Infection, Department of Internal Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

⁶Department of Internal Medicine, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia

*Correspondence Email: uzein_2000@yahoo.com; umar.zein@fk.uisu.ac.id

Abstract. Peritoneal tuberculosis may lead to delayed diagnosis because of the nonspecific features such as fever, abdominal distension, abdominal tenderness, ascites, and weight loss. Here, wereported a case of prolonged fever and abdominal pain which was due to peritoneal tuberculosis. Initial examinations including acomplete blood test and serologic tests did not lead to the diagnosis. A final diagnosis was made by abdominal CT-scan and laparoscopy combined with histopathological studies. Antituberculous medications provided a good clinical response in this patient.

1. Introduction

Peritoneal tuberculosis may lead to delayed diagnosis because of the nonspecific features such as fever, abdominal distension, abdominal tenderness, ascites, and weight loss.[1,2] Early diagnosis is very important to prevent morbidity and mortality. Prolonged fever and abdominal pain are common presenting symptoms. Abdominal CT-scan plays a major role in early diagnosis of this condition, but final diagnosis of peritoneal tuberculosis is based onthehistopathological study.[1]

Prolonged fever is significantly found more in children less than 5 years of age. However, one should suspect abdominal tuberculosis at any age group on prolonged fever followed by abdominal pain. Here, we describe a case of a 29-year-old man because of prolonged fever.

2. Case Report

A patient is a 29-year-old man Bataknes Indonesian, who came to the Tropical and Infectious Disease Clinic because of a two month of recurrent fever and a slightly enlarged abdominal wall. His weight had decreased by 6 kg. The weight was 58 kg. He was working in Miami Beach, USA when the symptoms occurred. He had been examined in the hospital.



Content from this work may be used under the terms of theCreative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

His past blood testing revealed normal complete blood count. The serologic testing including HIV, HCV, HBV, CMV, *Toxoplasma*, *Rubella*, *Epstein Barr*, and *Salmonella* was non-reactive. Urine screen was normal. There were no abnormalities in his chest X-ray and abdominal ultrasound. He had been treated with ceftriaxone, azithromycin, and moxifloxacin but the symptoms persisted.

On medical examination, the temperature was 38°C, the respiratory rate was 18 breaths per minute, the pulse 80 beats per minute, the blood pressure 130/80 mmHg. There was no lymph node enlargement. The abdomen was slightly enlarged and mild tenderness in the umbilical region, but peristalsis was normal.

The laboratory data results including complete blood counts, liver and kidney function tests were normal. The abdominal scan revealed aortic lymphadenopathy and peritonitis. Based on the clinical diagnosis of peritoneal tuberculosis, the anti-tuberculosis regimens of four drugs: rifampicin, isoniazid, ethambutol, and pyrazinamide were given.

The patient's conditions had been improved with no fever, increased appetite, and relieved abdominal pain. But his abdominal distension was more prominently associated with ascites. Thus, a diagnostic laparoscopy was performed. The results of laparoscopy revealed multiple small nodules in the peritoneum, intestines, and ascites (Figure 1). The ascitic fluid analysis was yellow and serous, LDH 288 U/L, protein 5.5 g/L, white cells count 180 cells/mm³, glucose level 79 mg/dL, pH 8.0, polymorphonuclear cells 50%, and mononuclear cells 50%. Ascitic fluid culture grew no bacteria. Ascitic fluid adenosine deaminase (ADA) was not measured. But histopathological examination of the peritoneal biopsy showed a specific chronic inflammation with lymphoid cells tubercles. The nodules contained epithelioid and Langhans giant cells and necrotic mass. Connective tissue stroma with inflammatory cell infiltration of lymphocytes was present.

The patient continued to take anti-tuberculous medications prescribed. On follow-up a month later, he reported increased appetite without fever. No abnormality was found on physical examination.




Figure 1. Laparoscopic appearance of peritoneal tuberculosis.

Discussion

The delay in the diagnosis of peritoneal tuberculosis resulted in prolonged fever in this patient, because of no specific features.[1] This condition is also a great mimicker of other abdominal conditions such as end-stage renal and liver diseases,[3] and ovarian carcinoma.[4]

Abdominal tuberculosis has an insidious course without any specific laboratory, radiological or clinical findings.[5] Peritoneal tuberculosis is a form of abdominal tuberculosis that involves intestinal tracts, liver, spleen, omentum, female genital tract, parietal and visceral peritoneum.[6] The non-specificity of clinical features leads to difficulty in its diagnosis.[5] Various investigative methods had been reported as the gold standard in earlier studies. However, there are difficulties in clinical practice. Thus, a prolonged fever and abdominal pain with weight loss should consider peritoneal tuberculosis in the differential diagnosis.

Clinical features of prolonged fever and ascites confirmed by histopathology studies of laparoscopy findings led to definitive diagnosis of peritoneal tuberculosis. Histopathological studies of the caseation in granulomas (nodules) are the hallmark of peritoneal tuberculosis.[1] Laparoscopic study and peritoneal biopsy are safety procedures in adult patients suspected of abdominal tuberculosis.[7] Bhargava et al. reported that laparoscopic findings by visual diagnosis were accurate in 95% of patients.[8] This examination enabled peritoneal biopsy to be made histologically on the basis of typical granulomas (nodules). Laparoscopy can be used for diagnosis of peritoneal tuberculosis with a sensitivity of 93% and a specificity of 98% when combined with histopathological studies.[9]

Most of the patients respond very well to the standard of antituberculous therapy.[1] Our patient's prominent symptom was prolonged fever without rigors and weight loss.[10] This unspecific features led to empiric therapy with antimicrobial regimes. In the present study, appropriate therapy with antituberculous medications led to the improvement of patient's condition. Although the symptoms had been improved after 1 month of standard antituberculous therapy, the treatment had to be continued for 6 months.[1]

4. Conclusion

Peritoneal tuberculosis may present with unspecific clinical symptoms and signs. Thus, it is regarded as a great mimicker of a variety of abdominal diseases. A prolonged fever followed by abdominal pain and weight loss should consider peritoneal tuberculosis. Once the definitive diagnosis has been made with combined laparoscopy and histopathological studies, the standard antituberculous drugs can be given.

References

- [1] Debi U, Ravisankar V, Prasad K K, Sinha S K and Sharma A K 2014 Abdominal tuberculosis of the gastrointestinal tract: Revisited *World J. Gastroenterol.* **20** 14831-40
- [2] Shah I and Uppuluri R 2010 Clinical profile of abdominal tuberculosis in children *Indian J. Med. Sci.* **64** 204-9 Available from: <http://www.indianjmedsci.org/text.asp?2010/64/5/204/98935> [Accessed: August 20, 2017]
- [3] Sokolina I, Reshetnikov M, Zyuzya Y, Guznov V and Sinitzjn M 2015 Computed tomography in abdominal tuberculosis in patients with HIV/tuberculosis co-infection *Eur. Soc. Radiol. Congress* DOI: 10.1594/ecr2015/C-224
- [4] Boss J D, Shah C T, Oluwole O and Sheagren J N 2012 TB peritonitis mistaken for ovarian carcinomatosis based on an elevated CA-125 *Case Report Med.* 1-3
- [5] Uygur-Bayramiçli O, Dabak G and Dabak R 2003 A clinical dilemma: abdominal tuberculosis *World J. Gastroenterol.* **9** 1098-101
- [6] Yazdani S, Sadeghi M, Alijanpour A and Naeimi-rad M 2016 A case report of peritoneal tuberculosis with multiple military peritoneal deposits mimicking advanced ovarian carcinoma *Caspian J. Intern. Med.* **7** 61-3
- [7] Erdogan D, Yıldız Y T, Boduroglu E C and Tanir N G 2013 An abdominal tuberculosis case mimicking an abdominal mass *Ann. Ped. Surg.* **9** 81-3
- [8] Bhargava D K, Shriniwas, Chopra P, Nijhawan S, Dasarathy S and Kushwaha A K 1992 Peritoneal tuberculosis: Laparoscopic patterns and its diagnostic accuracy *Am. J. Gastroenterol.* **87** 109-12
- [9] Rasheed S, Zinicola R, Watson D, Bajwa A and McDonald P J 2007 Intra-abdominal and gastrointestinal tuberculosis *Colorectal Dis.* **9** 773-83
- [10] Bernando J 2017 Clinical manifestations, diagnosis, and treatment of extrapulmonary and military tuberculosis Available from: <http://www.uptodate.com/contents/clinical-manifestations-diagnosis-and-treatment-of-extrapulmonary-and-military-tuberculosis> [Accessed: August 20, 2017]

Prolonged_Fever_in_Peritoneal_TB-compressed.pdf

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4