

- **April 26, 2020 (Important)**

Article No.: **102298-AJCS-ANSI**

Article Type: **Research Article**

**The article still has some reference issue. Kindly consider following comments and modify accordingly;**

- It is mandatory to provide active and trackable DOI or URL of all listed references. Author is advised to provide trackable active DOI or direct links highlighted References or replace the reference with suitable and trackable reference.
- There must be no duplication of any reference in Reference List. Each reference must be listed only once in Reference list. Reference No 16 is duplication of Reference No. 15. Author is advised to recheck all references and make sure there would be duplication of any reference.

## **ABA CONTENT OF PALM OIL SEEDLINGS (*Elaeis guineensis* Jacq) WITH VEDAGRO TREATMENT ON WATER STRESS**

<sup>1</sup>Bilter Sirait, <sup>2</sup>Agnes Imelda Manurung,  
<sup>3</sup>Ernitha Panjaitan and <sup>3</sup>Lince Siregar

<sup>1</sup>LLDikti Wil.I employed of Darma Agung University Medan 20153;  
<sup>2</sup>Department of Agriculture, Darma Agung University Medan 20153,  
<sup>3</sup>Department of Agriculture, Methodist Indonesia University, Medan 20132

**Corresponding Author: Bilter Sirait**  
**[dapejel.rait@yahoo.com](mailto:dapejel.rait@yahoo.com)**

**Conflict of interest: The authors have declared that no competing interest exists.**

### **Author's Contribution:**

Name of the co-author and e-mail ID	Types of contribution
Bilter Sirait. <a href="mailto:dapejel.rait@yahoo.com">dapejel.rait@yahoo.com</a>	-To Lead a discussion about journal target. -putting together a manuscript -double check the manuscript -report to the Asian Journal of Crop Science
Agnes Imelda Manurung. <a href="mailto:manurunghutabarat@gmail.com">manurunghutabarat@gmail.com</a>	-putting together a manuscript -submitting to internal reviewer -checking the manuscript with a grammarly machine
Ernitha Panjaitan. <a href="mailto:ernitha2005@yahoo.co.id">ernitha2005@yahoo.co.id</a>	-putting together a manuscript

	-checking the manuscript with a grammarly machine
LinceSiregar. <a href="mailto:meddy.siregar@yahoo.com">meddy.siregar@yahoo.com</a>	-putting together a manuscript

## ABSTRACT

**Background and Objective:** Indonesia is one of the chief producers of palm oil of the world market, but all production is collected at one place. This research will include later whether there is a correlation between seedlings that are more adaptable of water stress in nurseries with field planting. This study was aimed to know the response of the growth of palm oil seedlings and ABA content to Vedagro fertilizer treatment and water stress. **Materials and Methods:** The experiment was conducted in field experiment station of the Agricultural Faculty of DarmaAgung University Medan, Indonesia, with  $\pm$  28 meters above sea level. The research is done in May 2019 until August 2019. This research uses the Completely Randomized Block Design Factorial Method consisting of 2 treatment factors, that is Vedagroconsists of 3 levels, V1 = 5.0 g baby polybag<sup>-1</sup>, V2 = 10.0g and V3 = 15.0g baby polybag<sup>-1</sup> each, while water stress with 4 levels namely W0 = 0.2 litres baby polybag<sup>-1</sup>day<sup>-1</sup>; W1 = 0.2 litres baby polybag<sup>-1</sup>days<sup>-2</sup>; W2 = 0.2 litres baby polybag<sup>-1</sup>days<sup>-3</sup>; W3 = 0.2 litres baby polybag<sup>-1</sup>days<sup>-4</sup>. **Results:** The results showed that the higher the dose of Vedagro, the higher the growth of seedlings, on the contrary, the lack of water cause the growth of seedling decreased but the ABA content increased. **Conclusion:** Vedagro up to 15 g baby polybag<sup>-1</sup> increase significantly only to plant height; Water stress decreases significantly in plant growth but ABA content increase significantly 0.0022 ppm cm<sup>-2</sup> leaf area.

Keywords: *Vedagro*, *water stress*, *growth*, *seedlings*, *ABA*, *biodiversity*, *seedling*

## INTRODUCTION

Indonesia always adheres to the constitution of the Republic of Indonesia to participate for welfare of all the people and countries in the World, it is proven that Indonesia is one of the chief producers of oil palm has growth from 10.6 million ha in 2013 to 13.7 million ha by 2020<sup>1</sup> due to a ceaselessly growing demand<sup>2</sup>, eventhough Indonesia often receive negative issues against the palm oil industry such as the destruction of forests and biodiversity, food security and agrarian conflicts and natural resources.

Tenera oil palm which is a Simalungun variety of North Sumatra is a reliable variety that can be grown in various regions in the country. Description of the Simalungun variety shown that it can grow in various regions and has high productivity, although it is assumed that productivity will be different in line with differences in rainfall or water adequacy. Unfortunately, Simalungun varieties after being planted in several regions in Indonesia, then

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

companies or farmers harvesting, farmers collect and mix all fresh fruit bunches altogether means that all production is collected at one place so that they do not know the separation of each variety so the harvest is a collection of various varieties. This research will include later to solve the problem, one of them is whether there is a correlation between seedlings that are more adaptable of water stress in nurseries with field planting in various places in Indonesia where there are areas with low rainfall, moderate to areas of oil palm plantations with high rainfall. Morphological, physiological, and biochemical characterization needs to be carried out at later research, in previous research showed that the impact of water deficit are young leaves not opening, broken old leaf midrib until broken<sup>3</sup>.

Nowadays manure is very difficult to find and complicated, fertilizer Vedagro can replace it<sup>4</sup>. The need for water for oil palm seedlings in the pre-nursery is 0.2 litres baby polybag<sup>-1</sup>day<sup>-1</sup>, so the watering below 0.2 litres baby polybag<sup>-1</sup>day<sup>-1</sup> is considered to be water stresswhile in Malaysia watering 0.5 liters polybag<sup>-1</sup>day<sup>-1</sup> in the pre nursery because the polybag size is different<sup>5-7</sup>. From the results of previous studies, it is known that ABA is formed as a response to water deficits and it accumulates in all organs of the plant, is reported higher in leaves. In this regard, research needs to be done in the form of treatment of fertilizer Vedagro under conditions of water stress on the growth and development of oil palm seedlings (*Elaeisguineensis*Jacq). The purpose of this study is to know the growth of oil palm seedlings with Vedagro fertilizer treatment and ABA content due to lack of water.

## **MATERIALS AND METHODS**

### **Study area:**

The experimentwas conducted in the field experimental station of the the Agricultural Faculty of DarmaAgung University in Medan North Sumatera with ± 28 meters above sea level. This experiment uses Tenera oil palm seedlings variety of Simalungun which is obtained from the Indonesia Oil Palm Research Institute. Other material used is baby

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

polybag, planting medium, was carried out in a plastic house which is avoided of rainfall. The research is done in May 2019 until August 2019.

### **Methodology:**

The research procedure was started by filling baby polybag with planting media, then a week later planted oil palm seedlings and added with vedagro fertilizer according to treatment, then the data collection was adjusted to the response variable which was observed, once every two weeks, starting at the age of the seedlings 4 weeks to 12 weeks after planting. This experiment uses the Block Random Design Factorial Method consisting of 2 treatment factors. The first, Vedagro Fertilizer Dose consists of 3 levels, namely,  $V_1 = 5.0\text{g baby polybag}^{-1}$ ,  $V_2 = 10.0\text{g baby polybag}^{-1}$ ,  $V_3 = 15.0\text{g baby polybag}^{-1}$ , while the second, water stress consisting of 4 levels namely  $W_0 = 0.2\text{ litres baby polybag}^{-1}\text{day}^{-1}$ ;  $W_1 = 0.2\text{ litres baby polybag}^{-1}\text{days}^{-2}$ ;  $W_2 = 0.2\text{ litres baby polybag}^{-1}\text{days}^{-3}$ ;  $W_3 = 0.2\text{ litres baby polybag}^{-1}\text{days}^{-4}$ . The parameters were observed at ages 4-12 weeks after planting i.e. plant height, stem diameter, the number of leaves, leaf area, and the ABA content.

### **Statistical analysis:**

Statistical analysis using variance analysis, and if the treatment effect is significant, followed with the LSD test at the 5% level.

ABA content analysis is carried out following the procedure Laboratory of the Institute of Research and Development of Agriculture-Center for Research and Development of Agriculture Biotechnology and Genetic Resources of Agriculture, Bogor.

## **RESULTS**

Plant height. The results showed that plant height continued to increase with increasing plant age (Table 1). Although the effect of the combination treatment of Vedagro and Water Stress is not significant, however, the higher the dose of vedagro fertilizer the higher the plant height was found, on the other hand, the less water the more depressed the

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

growth of seedlings, Watering 0.2 litres baby polybag<sup>-1</sup>days<sup>-4</sup> significantly reduces the height of oil palm seedlings as many 7.81 cms at 12 weeks after planting.

**Table 1:** Effect of Vedagro Fertilizer and Water Stress Treatment to Plant Height of Palm Oil Seedlings at Ages 4 to 12 Weeks After Planting (WAP)

Treatment	Plant height (cm)				
	4 WAP	6 WAP	8 WAP	10 WAP	12 WAP
V <sub>1</sub> =5 g baby polybag <sup>-1</sup>	9.12	12.66	15.35	18.36a	21.54a
V <sub>2</sub> =10 g baby polybag <sup>-1</sup>	9.84	13.51	16.22	19.18ab	22.61b
V <sub>3</sub> =15 g baby polybag <sup>-1</sup>	9.39	12.99	15.79	19.91b	23.30c
W <sub>0</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-1</sup>	9.60	13.28	16.69	20.50 a	23.96a
W <sub>1</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-2</sup>	9.50	13.03	15.79	19.92a	23.47a
W <sub>2</sub> =0.2 liters baby polybag <sup>-1</sup> days <sup>-3</sup>	9.53	13.07	15.19	18.19 ab	22.35ab
W <sub>3</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-4</sup>	9.27	12.82	13.09	15.99 b	16.15b

Note: Numbers followed by the same letter in the same column and the same treatment are not different significantly with LSD test at the 5% level

Stem Diameter. The diameter of the plants continues to increase with increasing ages (Table 2). The pattern is the same, the higher the dose of fertilizer the higher the stem diameter was found, on the other hand, the less water the more depressed stem diameter of seedlings, Watering 0.2 litres baby polybag<sup>-1</sup>days<sup>-4</sup> significantly reduces stem diameter of oil palm seedlings as many 3,79 cms at 12 weeks after planting.

**Table 2:**Effect of Vedagroand Water Stress Treatment to Stem Diameter of Palm Oil Seedling at Ages 4 to 12 Weeks After Planting (WAP)

Treatment	Stem Diameter (cm)				
	4 WAP	6 WAP	8 WAP	10 WAP	12 WAP
V <sub>1</sub> =5 g baby polybag <sup>-1</sup>	3.25	4.33	4.59	4.92	5.52
V <sub>2</sub> =10 g baby polybag <sup>-1</sup>	3.19	4.37	4.56	5.06	5.74
V <sub>3</sub> =15 g baby polybag <sup>-1</sup>	3.30	4.41	4.80	5.45	6.06
W <sub>0</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-1</sup>	3.29	4.48	4.95	5.54	6.20a
W <sub>1</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-2</sup>	3.26	4.31	4.71	5.17	5.91ab
W <sub>2</sub> =0.2 liter baby polybag <sup>-1</sup> days <sup>-3</sup>	3.23	4.44	4.54	5.02	5.56a
W <sub>3</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-4</sup>	2.20	2.24	2.39	2.40	2.41b

Note: Numbers followed by the same letter in the same column are not different significantly with LSD test at the 5% level

The number of leaves continues to increase with increasing age (Table 3) although the effect of vedagro fertilizer is not significant. The effect of water stress is also not significant on the number of leaves.

**Table 3:**Effect of Vedagro and Water Stress Treatment to Number of Leaves of Palm Oil Seedling at Ages 4 to 12 Weeks After Planting (WAP)

Treatment	The Number of Leaves				
	4 WAP	6 WAP	8 WAP	10 WAP	12 WAP
V <sub>1</sub> =5 g baby polybag <sup>-1</sup>	2.64	3.62	4.59a	5.59	6.03
V <sub>2</sub> =10 g baby polybag <sup>-1</sup>	2.67	3.67	4.62b	5.70	6.31
V <sub>3</sub> =15 g baby polybag <sup>-1</sup>	2.73	3.73	4.73c	5.89	6.39
W <sub>0</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-1</sup>	2.71	3.71	4.71b	5.89	6.60
W <sub>1</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-2</sup>	2.71	3.71	4.67b	5.71	6.52
W <sub>2</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-3</sup>	2.71	3.71	4.71b	5.82	6.11
W <sub>3</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-4</sup>	2.60	3.56	4.49a	5.49	5.74

Leaf area due to Vedagro fertilizer dose treatment and water treatment at ages 12 weeks after planting (WAP) shown in Table 4. The effect of Vedagro fertilizer and the combination treatment of vedagro and water stress is not significant. Watering 0.2 litres baby polybag<sup>-1</sup>days<sup>-4</sup> significantly reduces the leaf area of oil palm seedlings as many 20,82 cm<sup>2</sup> at 12 weeks after planting.

**Table 4:** Leaf area plant due to Vedagro Fertilizer Dose Treatment and Water Stress Treatment at Ages 12 Weeks After Planting (WAP)

Treatment	W <sub>0</sub> 0.2litres baby polybag <sup>-1</sup> days <sup>-1</sup>	W <sub>1</sub> 0.2 litres baby polybag <sup>-1</sup> days <sup>-2</sup>	W <sub>2</sub> 0.2 litres baby polybag <sup>-1</sup> days <sup>-3</sup>	W <sub>3</sub> 0.2 litres baby polybag <sup>-1</sup> days <sup>-4</sup>	Average
V <sub>1</sub> =5 g baby polybag <sup>-1</sup>	32.01	31.14	18.17	11.10	23.11
V <sub>2</sub> =10 g baby polybag <sup>-1</sup>	32.17	31.46	19.23	11.27	23.53
V <sub>3</sub> =15 g baby polybag <sup>-1</sup>	32.97	32.60	23.50	12.30	25.34
Average	32.38a	31.73a	20.30b	11.56c	23.99

Note: Numbers followed by the same letter in the same row are not different significantly with LSD test at the 5% level

The observational matrix at age 12 WAP with water stress treatment is shown in Table 5. Increasingly lack water, all parameters observed were significantly reduced except for the number of leaves is not significant. Increasingly lack water, ABA content increased in leaf. The regression equation of water stress with ABA content is shown in Figure 1. Figure 1 shows that Increasingly lack water in palm oil seedlings at ages 12 WAP then the ABA content increased,  $Y = 1.82 + 0.71 X$ ,  $r = 0.99$ . This can be interpreted in agronomic approach that each decrease of 1 liter in watering will increase the ABA content = 0.90 ppm.

**Table 5:**The Observational Matrix at Age 12 WAP with Water Stress Treatment

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

Treatment	Plant Height(cm)	Stem Diameter (cm)	The Nnumber of Leaves	Leaf Area(cm <sup>2</sup> )	ABA (ppm)
W <sub>0</sub> = 0.2 litres baby polybag <sup>-1</sup> day <sup>-1</sup>	23.96a	6.20a	6.60ns	32.38a	2.52b
W <sub>1</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-2</sup>	23.47ab	5.91 ab	6.52 ns	31.73a	3.46ab
W <sub>2</sub> =0.2 liter baby polybag <sup>-1</sup> days <sup>-3</sup>	22.35ab	5.56 a	6.11 ns	20.30b	3.53ab
W <sub>3</sub> =0.2 litres baby polybag <sup>-1</sup> days <sup>-4</sup>	16.15b	2.41 b	5.74 ns	11.56c	4.86a

Note: Numbers followed by the same letter in the same column are not different significantly with the LSD test at the 5% level




Figure 1. Relationship between Water Stress and ABA Content at Ages 12 WAP

## DISCUSSION

Fertilizer can add nutrients in the soil, so that plant growth increases with the availability of nutrients. Fertilizers that are put into the soil will be decomposed by microorganisms and nutrients released and become available or can be absorbed by plant roots. Overall, Vedagro fertilizer can increase the growth of oil palm seedlings. This is alleged because it can supply nutrients, improve the physical, chemical and biological properties of the soil, in turn, the soil can provide nutrients in a balanced amount. These

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

results are not much different from those obtained previously on cassava that Vedagro fertilizer increased tuber yield about 79 %<sup>8</sup>. Application of Vedagro fertilizer at a dose of 5 g, 10 g, 15 g, and 20 g / baby polybag significantly affects plant height, leaf area, total dry weight of plants. Vedagro fertilizer containing macro nutrients such as N: 11-12%, K<sub>2</sub>O: 4.5-6.0%, P<sub>2</sub>O<sub>5</sub>: 0.4-0.6%, Ca: 1.1%, Mg: 1.9- 2.2%, and micro nutrients consisting of Fe, Mn, Cu, Zn, B and Mo are suitable for increasing growth of oil palm seedlings. Vedagro fertilizers also contain aspartic acid, glutamic acid, Alanine, Valine, Isoleucine, Leucine, Tyrosine, Lysine, and Arginine<sup>4</sup>.

Each plant has a different sensitivity to water stress, but tillering and vegetative growth stage were the target of many studies because they are important growth phases from a crop production perspective<sup>9</sup>. Water stress decreases plant growth, and this is very closely related to plant metabolism, ABA is produced in large quantities as a mechanism of adaptation of plants to water scarcity/drought or dehydration<sup>9-11</sup>. ABA leads to feedback inhibition of photosynthesis with the aid of carbohydrate accumulation and decline in the concentration of photosynthetic enzymes which result in the sequencing of photosynthesis arrest, automatically, growth of plants also arrest. Water stress-causing osmotic stresses and cellular dehydration, and water-stressed vegetation have more concentration of ABA than the good-watered crops, which restricts growth predominantly inhibit shoot growth<sup>9-12</sup> and supplying 75% water from soil field capacity on oil palm still possible<sup>13</sup>, oil palm plant with water scarcity significantly reduces growth and oil yield but elevated ABA and sitokinin content as a typical osmoprotectant to maintain cell stability during cell dehydration which is the common response of plant to drought stress<sup>14-15</sup>.

Implication of this research is to facilitate the selection and application of fertilizer for oil palm seedlings, means that the higher the vedagro fertilizer as classified as organic fertilizer, the higher growth of oil palm seedlings is. As a recommendation of this study is



Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

effect of vedagro fertilizer would be significant on oil palm seedlings at high doses of 15 g baby polybag<sup>-1</sup>, while the watering should be according to the suggestion of 0.2 liters of baby polybag<sup>-1</sup> day<sup>-1</sup> water. Limitation of this study, has not been tried in multi-location dry areas in Indonesia and needs to increase response variables.

## **CONCLUSION**

Vedagro Fertilizer up to 15 g baby polybag<sup>-1</sup> increase significantly plant height, but not significant effect to stem diameter, the number of leaves and leaf area of oil palm seedlings in the pre-nursery. Water stress decreases significantly in plant growth, namely plant height, stem diameter, number of leaves, leaf area of oil palm seedlings but ABA content increase significantly. Water stress can increase the ABA content as many 0.0022 ppm cm<sup>-2</sup> leaf area. The interaction of doses of Vedagro fertilizer and water stress has no significant effect on plant height, stem diameter, number of leaves and leaf area of oil palm seedlings in the pre-nursery.

## **SIGNIFICANCE STATEMENTS**

This Study discover the effect of Vedagro fertilizer to produce better growth of oil palm seedlings and the effect of water stress on seedling growth. This study will help researcher, farmers, seed industry to uncover the critical areas on nursery in various places in Indonesia. Thus, a new theory is found that on water stress can be predicted ABA content formed, may be arrived at.

## **ACKNOWLEDGMENT**

We express our gratitude to the Board of Darma Agung University Foundation and Rector of Darma Agung University for their help both materially and morally. Thanks are also extended to Sitorus Lature, Gunania, Ruth Maria, Klintan Sinurat, Epianus Ndraha, Feri Sipahutar, Irwan Jaya, Fidelis Gea, Jery Frans, Sentrius Bu'ulolo for their technical assistance.

## REFERENCES

- ~~1. Ferrianta, Y. (2017). Issue and Solution Palm Oil Industry in Indonesia. Agriculture Research and Technology 5:1 2. Available from: <https://juniperpublisher.com/artoaj/pdf/ARTOAJ.MS.ID.555656.pdf>~~
1. Shahputra and Z.Zen, 2018. Positive and negative impacts of oil palm expansion in Indonesia and the prospect to achieve sustainable palm oil. IOP Conf.Series: Earth and Environmental Science 122 (2018)012008. Doi:10.1088/1755-1315/122/1/012008. URL:<https://iopscience.iop.org/article/10.1088/1755-1315/122/1/012008/pdf>
2. Arunachalam, 2012. 3 - Oil palm. Genomics of cultivated palms, p.29-48. Elsevier. Available from: <https://doi.org/10.1016/B978-0-12-387736-9.00003-0>. URL: <https://www.sciencedirect.com/science/article/pii/B9780123877369000030?via%3Dihub>
3. Abd.Gafur, M. and E.T.Susila Putra, 2019.Effect of drought stress in physiological oil palm seedling (*Elaeisguineensis*Jacq.) using cacium application. Asian Journal of Biological Sciences, 12(3):550-556. DOI:[10.3923/ajbs.2019.550.556](https://doi.org/10.3923/ajbs.2019.550.556). URL:<https://scialert.net/abstract/?doi=ajbs.2019.550.556>
4. Linn, K.Z. and P.P. Myint, 2018. Study on the effectiveness of natural organic fertilizers on Cassava (*Manihotesculenta*Crantz.) cultivation. Asian J. of Soil Sci. and Plant Nutrition 3(3): 1-10, 2018; Article no.AJSSPN.42912 ISSN: 2456-9682. DOI: 10.9734/AJSSPN/2018/42912.URL:<http://journalajsspn.com/index.php/AJSSPN/article/download/999/1398/>
5. Osakabe Y., K. Osakabe, K. Shinozaki and Lam Son P. Tran, 2014. Response of plants to water stress. Front.Plant Sci.,13 March 2014. <https://doi.org/10.3389/fpls.2014.00086>. URL:<https://www.frontiersin.org/articles/10.3389/fpls.2014.00086/full>
6. Muhamad, H., V. Subramaniam, Z.Hashim, N.S.K.Khairuddin and C.Y.May., 2014. Water footprint: part 1 - production of oil palm seedlings in Peninsular Malaysia. Journal of Oil Palm Research, 26 (4): 273-281. Available from: [https://www.researchgate.net/publication/279335013\\_Water\\_footprint\\_Part\\_1\\_-\\_Production\\_of\\_oil\\_palm\\_seedlings\\_in\\_peninsular\\_Malaysia](https://www.researchgate.net/publication/279335013_Water_footprint_Part_1_-_Production_of_oil_palm_seedlings_in_peninsular_Malaysia)
7. Santosa, E., I. M. Stefano, A.G.Tarigan, A.on established Wachjar, S.Zaman and H. Augusta, 2018. Tree-based water footprint assessment oil palm plantation in North Sumatera, Indonesia. J. Agron. Indonesia, 46(1):111-118. DOI: <https://dx.doi.org/10.24831/jai.v46i1.13665>. ISSN 2085-2916 e-ISSN 2337-3652.<http://jai.ipb.ac.id>

Mastur, 2016. Physiological Responses of Sugar Cane to Drought. Industrial Tobacco, Fiber and Oil Plant Bulletin 8 (2), October 2016: 98-111. <http://ejurnal.litbang.pertanian.go.id/bultas->issue->download->pdf-2>

8. Linn, K. Z. and P. P. Myint, 2018. Study on the effectiveness of natural organic fertilizers on cassava (*Manihotesculenta*Crantz.) Cultivation. Asian Journal of Soil Science and Plant Nutrition 3(3): 1-10, 2018; Article no.AJSSPN.42912 ISSN: 2456-9682. DOI: 10.9734/AJSSPN/2018/42912 URL: [https://www.researchgate.net/publication/327264921\\_Study\\_on\\_the\\_Effectiveness\\_of\\_Natural\\_Organic\\_Fertilizers\\_on\\_Cassava\\_Manihot\\_esculenta\\_Crantz\\_Cultivation](https://www.researchgate.net/publication/327264921_Study_on_the_Effectiveness_of_Natural_Organic_Fertilizers_on_Cassava_Manihot_esculenta_Crantz_Cultivation)
  9. Ferreira, T.H.S, M.S.Tsunada, D.Bassi, P.Araujo, L.Mattiello, G.V. Guidelli, G.L.Righetto, P.Lakshmanan and M.Menossi, 2017. Sugarcane water stress tolerance mechanisms and its implications on developing biotechnology solutions.[Front Plant Sci.](#) 2017; 8: 1077. Published online 2017 Jun 23. doi: [10.3389/fpls.2017.01077](https://doi.org/10.3389/fpls.2017.01077)
  10. Chaves, M.M., Maroco, J.P., and Pereira, J.S., 2003. Understanding plant responses to drought from genes to the whole plant. *Functional Plant Biology*, 30. 239-264. [www.publish.csiro.au/journals/fpb](http://www.publish.csiro.au/journals/fpb). DOI: [10.1071/FP02076](https://doi.org/10.1071/FP02076).
- Hanafi, Sirait, S.M., Purwanti E., Irawan, C., Rochaeni, H., Rosita, T., and Purba, J.A., 2017. Comparison of Abscisic Acid Hormone on the leaves of Palm Oil Seeds with Watered and not watered. *Journal of Pharmacognosy and Phytochemistry*. [www.phytojournal.com](http://www.phytojournal.com). E-ISSN:2278-4136 P-ISSN:2349-8234
11. Hanafi, S. M. Sirait, Purwanti, E., Irawan., C., Rochaeni, H., Rosita, T., and Purba, J. A., 2018. Comparison of abscisic acid hormone on the leaves of palm oil seeds with watered and not watered. *Journal of Pharmacognosy and Phytochemistry* 2018; 7(1):410-413. Available from:<http://www.phytojournal.com/archives/2018/vol7issue1/PartF/6-6-323-936.pdf>
  12. Salazar, C., C. Hernandez, and M.T. Pino, 2015. Plant water stress; associations between ethylene and abscisic acid response. *Chilean Journal of Agricultural Research* 75 (suppl.1).[https://scielo.conicyt.cl/scielo.php?script=sci\\_arttext&pid=S0718-8392015000300008](https://scielo.conicyt.cl/scielo.php?script=sci_arttext&pid=S0718-8392015000300008).versión On-line ISSN 0718-5839. Available from:[https://scielo.conicyt.cl/scielo.php?script=sci\\_arttext&pid=S0718-58392015000300008](https://scielo.conicyt.cl/scielo.php?script=sci_arttext&pid=S0718-58392015000300008)
  13. Najihah, T.S., M. H.Ibrahim, A.A. Razak, 2019. Effect of water stress on the growth, physiology and biochemical properties of oil palm seedlings. [AIMS Agriculture and Food](#), 2019, 4(4): 854-868. doi: [10.3934/agrfood.2019.4.854](https://doi.org/10.3934/agrfood.2019.4.854).
  14. Legros, S., Mialet-Serra, I., Caliman, J.P., Siregar, F.A., Clement-Vidal, A., and Dingkuhn, M. (2009). Phenology and growth adjustmants of oil palm (*Elaeisguineensis*Jacq) to photoperiod and climate variability. *Annals of Botany* 104:1171-1182. doi:10.1093/aob/mcp214, Available from:[www.aob.oxfordjournals.org](http://www.aob.oxfordjournals.org)

Growth and ABA content on palm oil seedlings (*Elaeisguineensis*Jacq.)

15. Duangpan, S., Sujitto, S and Eksomtramage, T., 2017. Genotypic variation in proline accumulation during sequential drought and rewatering in response to drought preconditioning. International Journal of Agricultural Technology 2017. Vol.13 (6): 927-940. Available from: <http://www.ijat-aatsea.com> ISSN 1686-9141

~~Cao *et al*, 2011; Carr, 2011 in Duangpan, S., Sujitto, S and Eksomtramage, 2017. Genotypic variation in Proline Accumulation during Sequencing Drought and Rewatering in Response to Drought Preconditioning International Journal of Agricultural Technology 2017, 13 (6): 927-940. <http://www.ijat-aatsea.com>~~

- ~~16. Duangpan, S., Sujitto, S and Eksomtramage, T., 2017. Genotypic Variation in Proline Accumulation during Sequential Drought and Rewatering in Response to Drought Preconditioning. International Journal of Agricultural Technology 2017. Vol.13 (6):927-940. Available from: <http://www.ijat-aatsea.com> ISSN 2630-0192.~~



# Plagiarism Checker X Originality Report

**Similarity Found: 13%**

Date: Saturday, April 04, 2020

Statistics: 400 words Plagiarized / 3078 Total words

Remarks: Low Plagiarism Detected - Your Document needs Optional Improvement.

---

ABA CONTENT OF PALM OIL SEEDLINGS (*Elaeis guineensis* Jacq) WITH VEDAGRO TREATMENT ON WATER STRESS 1Bilter Sirait\*,2Agnes Imelda Manurung, 3Ernitha Panjaitan and 3Lince Siregar 1LLDikti Wil.I employed of Darma Agung University Medan 20153; 2Department of Agriculture, Darma Agung University Medan 20153, 3Department of Agriculture, Methodist Indonesia University, Medan 20132  
\*Corresponding Author: +628126559856 dapejel.rait@yahoo.com  
ABSTRACT  
Background and Objective: Indonesia is one of the chief producers of palm oil of the world market, but all production is collected at one place.

This research will include later whether there is a correlation between seedlings that are more adaptable of water stress in nurseries with field planting. This study aims to know the response of the growth of palm oil seedlings and ABA content to Vedagro fertilizer treatment and water stress. Materials and Methods: The experiment was conducted in field experiment station of the Agricultural Faculty of Darma Agung University Medan, Indonesia, with  $\pm 28$  meters above sea level. The research is done in May 2019 until August 2019.

This research uses the Completely Randomized Block Design Factorial Method consisting of 2 treatment factors, that is Vedagro consists of 3 levels, V1 = 5.0 g baby polybag-1, V2 = 10.0 g and V3 = 15.0 g baby polybag-1 each, while water stress with 4 levels namely W0 = 0.2 litres baby polybag-1day-1; W1 = 0.2 litres baby polybag-1days-2; W2 = 0.2 litres baby polybag-1days-3; W3 = 0.2 litres baby polybag-1days-4.

Results: The results showed that the higher the dose of Vedagro, the higher the growth of seedlings, on the contrary, the lack of water cause the growth of seedling decreased

but the ABA content increased. Conclusion: Vedagro up to 15 g baby polybag-1 increase significantly only to plant height; Water stress decreases significantly in plant growth but ABA content increase significantly 0.0022 ppm cm<sup>-2</sup> leaf area.

Keywords: fertilizer, water stress, growth, and ABA content INTRODUCTION Indonesia always adheres to the constitution of the Republic of Indonesia to participate for welfare of all the people and countries in the World, it is proven that Indonesia is one of the chief producers of oil has already mastered 37% of the world market<sup>1</sup> due to a ceaselessly growing demand<sup>2</sup>, even though Indonesia often receive negative issues against the palm oil industry such as the destruction of forests and biodiversity, food security and agrarian conflicts and natural resources.

Tenera oil palm which is a Simalungun variety of North Sumatra is a reliable variety that can be grown in various regions in the country. Description of the Simalungun variety shown that it can grow in various regions and has high productivity, although it is assumed that productivity will be different in line with differences in rainfall or water adequacy.

Unfortunately, Simalungun varieties after being planted in several regions in Indonesia, then companies or farmers harvesting, farmers collect and mix all fresh fruit bunches altogether means that all production is collected at one place so that they do not know the separation of each variety so the harvest is a collection of various varieties.

This research will include later to solve the problem, one of them is whether there is a correlation between seedlings that are more adaptable of water stress in nurseries with field planting in various places in Indonesia where there are areas with low rainfall, moderate to areas of oil palm plantations with high rainfall. Morphological, physiological, and biochemical characterization needs to be carried out at later research, in previous research showed that the impact of water deficit are young leaves not opening, broken old leaf midrib until broken<sup>3</sup>. Nowadays manure is very difficult to find and complicated, fertilizer Vedagro can replace it<sup>4</sup>.

The need for water for oil palm seedlings in the pre-nursery is 0.2 litres baby polybag-1day-1, so the watering below 0.2 litres baby polybag-1day-1 is considered to be water stress<sup>5,6,7</sup>. From the results of previous studies, it is known that ABA is formed as a response to water deficits and it accumulates in all organs of the plant, is reported higher in leaves.

In this regard, research needs to be done in the form of treatment of fertilizer Vedagro under conditions of water stress on the growth and development of oil palm seedlings

(*Elaeis quineensis* Jacq). The purpose of this study is to know the growth of oil palm seedlings with Vedagro fertilizer treatment and ABA content due to lack of water.

**MATERIALS AND METHODS** The experiment was conducted in the field experimental station of the the Agricultural Faculty of Darma Agung University in Medan North Sumatera with  $\pm 28$  meters above sea level. This experiment uses Tenera oil palm seedlings variety of Simalungun which is obtained from the Indonesia Oil Palm Research Institute. Other material used is baby polybag, planting medium, was carried out in a plastic house which is avoided of rainfall.

The research is done in May 2019 until August 2019. The research procedure was started by filling baby polybag with planting media, then a week later planted oil palm seedlings and added with vedagro fertilizer according to treatment, then the data collection was adjusted to the response variable which was observed, once every two weeks, starting at the age of the seedlings 4 weeks to 12 weeks after planting. This experiment uses the Block Random Design Factorial Method consisting of 2 treatment factors.

The first, Vedagro Fertilizer Dose consists of 3 levels, namely, V1 = 5.0 g baby polybag-1, V2 = 10.0 g baby polybag-1, V3 = 15.0 g baby polybag-1, while the second, water stress consisting of 4 levels namely W0 = 0.2 litres baby polybag-1day-1; W1 = 0.2 litres baby polybag-1days-2; W2 = 0.2 litres baby polybag-1days-3; W3 = 0.2 litres baby polybag-1days-4. The parameters were observed at ages 4-12 weeks after planting i.e.

plant height, stem diameter, the number of leaves, leaf area, and the ABA content. ABA content analysis is carried out following the procedure Laboratory of the Institute of Research and Development of Agriculture-Center for Research and Development of Agriculture Biotechnology and Genetic Resources of Agriculture, Bogor. RESULTS Plant height.

The results showed that plant height continued to increase with increasing plant age (Table 1). Although the effect of the combination treatment of Vedagro and Water Stress is not significant, however, the higher the dose of vedagro fertilizer the higher the plant height was found, on the other hand, the less water the more depressed the growth of seedlings, Watering 0.2

litres baby polybag-1days-4 significantly reduces the height of oil palm seedlings as many 7.81 cms at 12 weeks after planting. Table 1. Plant Height of Palm Oil Seedlings due to Vedagro Fertilizer Treatment and Water Stress Treatment at Ages 4, 6, 8, 10 and 12 Weeks After Planting (WAP)

Treatment	Plant height (cm)	_ _ _4 WAP	_6 WAP	_8 WAP
-----------	-------------------	------------	--------	--------

\_10 WAP \_12 WAP \_\_V1 =5 g baby polybag-1 \_9.12 \_12.66 \_15.35 \_18.36a \_21.54a \_\_V2 =10 g baby polybag-1 \_9.84 \_13.51 \_16.22 \_19.18ab \_22.61b \_\_V3=15 g baby polybag-1 \_9.39 \_12.99 \_15.79 \_19.91b \_23.30c \_\_W0=0.2 litres baby polybag-1days-1 \_9.60 \_13.28 \_16.69 \_20.50 a \_23.96a \_\_W1=0.2 litres baby polybag-1days-2 \_9.50 \_13.03 \_15.79 \_19.92a \_23.47a \_\_W2=0.2 liters baby polybag-1days-3 \_9.53 \_13.07 \_15.19 \_18.19 ab \_22.35ab \_\_W3=0.2

litres baby polybag-1days-4 \_9.27 \_12.82 \_13.09 \_15.99 b \_16.15b \_\_Note: Numbers followed by the same letter in the same column and the same treatment are not different significantly with LSD test at the 5% level Stem Diameter. The diameter of the plants continues to increase with increasing ages (Table 2).

The pattern is the same, the higher the dose of fertilizer the higher the stem diameter was found, on the other hand, the less water the more depressed stem diameter of seedlings, Watering 0.2 litres baby polybag-1days-4 significantly reduces stem diameter of oil palm seedlings as many 3,79 cms at 12 weeks after planting. Table 2.

Stem Diameter of Palm Oil Seedling Plant due to Vedagro Fertilizer Dose Treatment and Water Stress Treatment at Ages 4, 6, 8, 10 and 12 Weeks After Planting (WAP) Treatment \_Stem Diameter (cm) \_\_4 WAP \_6 WAP \_8 WAP \_10 WAP \_12 WAP \_\_V1 =5 g baby polybag-1 \_3.25 \_4.33 \_4.59 \_4.92 \_5.52 \_\_V2 =10 g baby polybag-1 \_3.19 \_4.37 \_4.56 \_5.06 \_5.74 \_\_V3=15 g baby polybag-1 \_3.30 \_4.41 \_4.80 \_5.45 \_6.06 \_\_W0=0.2 litres baby polybag-1days-1 \_3.29 \_4.48 \_4.95 \_5.54 \_6.20a \_\_W1=0.2

litres baby polybag-1days-2 \_3.26 \_4.31 \_4.71 \_5.17 \_5.91ab \_\_W2=0.2 liter baby polybag-1days-3 \_3.23 \_4.44 \_4.54 \_5.02 \_5.56a \_\_W3=0.2 litres baby polybag-1days-4 \_2.20 \_2.24 \_2.39 \_2.40 \_2.41b \_\_Note: Numbers followed by the same letter in the same column are not different significantly with LSD test at the 5% level The number of leaves continues to increase with increasing age (Table 3) although the effect of vedagro fertilizer is not significant. The effect of water stress is also not significant on the number of leaves.

Table 3. The Number of Leaves of Palm Oil Seedling Plant due to Vedagro Fertilizer Dose Treatment and Water Stress Treatment at Ages 4, 6, 8, 10 and 12 Weeks After Planting (WAP) Treatment \_The Number of Leaves \_\_4 WAP \_6 WAP \_8 WAP \_10 WAP \_12 WAP \_\_V1 =5 g baby polybag-1 \_2.64 \_3.62 \_4.59a \_5.59 \_6.03 \_\_V2 =10 g baby polybag-1 \_2.67 \_3.67 \_4.62b \_5.70 \_6.31 \_\_V3=15 g baby polybag-1 \_2.73 \_3.73 \_4.73c \_5.89 \_6.39 \_\_W0= 0.2 litres baby polybag-1days-1 \_2.71 \_3.71 \_4.71b \_5.89 \_6.60 \_\_W1=0.2 litres baby polybag-1days-2 \_2.71 \_3.71 \_4.67b \_5.71 \_6.52 \_\_W2=0.2 litres baby polybag-1days-3 \_2.71 \_3.71 \_4.71b \_5.82 \_6.11 \_\_W3=0.2



litres baby polybag-1days-4 2.60 3.56 4.49a 5.49 5.74 Leaf area due to Vedagro fertilizer dose treatment and water treatment at ages 12 weeks after planting (WAP) shown in Table 6. The effect of Vedagro fertilizer and the combination treatment of vedagro and water stress is not significant. Watering 0.2

litres baby polybag-1days-4 significantly reduces the leaf area of oil palm seedlings as many 20,82 cm<sup>2</sup> at 12 weeks after planting. Table 6. Leaf area plant due to Vedagro Fertilizer Dose Treatment and Water Stress Treatment at Ages 12 Weeks After Planting (WAP) Treatment W0 W1 W2 W3 Average 0.2litres baby polybag-1days-1 0.2 litres baby polybag-1days-2 0.2 litres baby polybag-1days-3 0.2 litres baby polybag-1days-4 V1 =5 g baby polybag-1 32.01 31.14 18.17 11.10 23.11 V2 =10 g baby polybag-1 32.17 31.46 19.23 11.27 23.53 V3=15 g baby polybag-1 32.97 32.60 23.50 12.30 25.34 Average 32.38a 31.73a 20.30b 11.56c 23.99 Note: Numbers followed by the same letter in the same row are not different significantly with LSD test at the 5% level The observational matrix at age 12 WAP with water stress treatment is shown in Table 7.

Increasingly lack water, all parameters observed were significantly reduced except for the number of leaves is not significant. Increasingly lack water, ABA content increased in leaf. The regression equation of water stress with ABA content is shown in Figure 1. Figure 1 shows that Increasingly lack water in palm oil seedlings at ages 12 WAP then the ABA content increased,  $Y = 1.82 + 0.71 X$ ,  $r = 0.99$ .

This can be interpreted in agronomic approach that each decrease of 1 liter in watering will increase the ABA content = 0.90 ppm. Table 7. The Observational Matrix at Age 12 WAP with Water Stress Treatment Treatment Plant Height (cm) Stem Diameter (cm) The Number of Leaves Leaf Area(cm<sup>2</sup>) ABA (ppm) W0= 0.2 litres baby polybag-1day-1 23.96a 6.20a 6.60ns 32.38a 2.52b W1=0.2 litres baby polybag-1days-2 23.47ab 5.91 ab 6.52 ns 31.73a 3.46ab W2=0.2

liter baby polybag-1days-3 22.35ab 5.56 a 6.11 ns 20.30b 3.53ab W3=0.2 litres baby polybag-1days-4 16.15b 2.41 b 5.74 ns 11.56c 4.86a Note: Numbers followed by the same letter in the same column are not different significantly with the LSD test at the 5% level / Figure 1. Relationship Between Water Stress and ABA Content at Ages 12 WAP DISCUSSION Fertilizer can add nutrients in the soil, so that plant growth increases with the availability of nutrients.

Fertilizers that are put into the soil will be decomposed by microorganisms and nutrients released and become available or can be absorbed by plant roots. Overall, Vedagro

fertilizer can increase the growth of oil palm seedlings. This is alleged because it can supply nutrients, improve the physical, chemical and biological properties of the soil, in turn, the soil can provide nutrients in a balanced amount.

These results are not much different from those obtained by that the application of Vedagro fertilizer at a dose of 5 g, 10 g, 15 g, and 20 g / baby polybag significantly affects plant height, leaf area, total dry weight of plants. Vedagro fertilizer containing macro nutrients such as N: 11-12%, K<sub>2</sub>O: 4.5-6.0%, P<sub>2</sub>O<sub>5</sub>: 0.4-0.6%, Ca: 1.1%, Mg: 1.9-2.2%, and micro nutrients consisting of Fe, Mn, Cu, Zn, B and Mo are suitable for increasing growth of oil palm seedlings.

Vedagro fertilizers also contain aspartic acid, glutamic acid, Alanine, Valine, Isoleucine, Leucine, Tyrosine, Lysine, and Arginine<sup>4</sup>. Each plant has a different sensitivity to water stress, but tillering and grant vegetative growth stage were the target of many studies because they are important growth phases from a crop production perspective<sup>9</sup>.

Water stress decreases plant growth, and this is very closely related to plant metabolism, ABA is produced in large quantities as a mechanism of adaptation of plants to water scarcity/drought or dehydration<sup>9,10,11</sup>. ABA leads to feedback inhibition of photosynthesis with the aid of carbohydrate accumulation and decline in the concentration of photosynthetic enzymes which result in the sequencing of photosynthesis arrest, automatically, growth of plants also arrest.

Water stress-causing osmotic stresses and cellular dehydration, and water-stressed vegetation have more concentration of ABA than the good-watered crops, which restricts growth predominantly inhibit shoot growth<sup>9,10,11,12</sup> and supplying 75% water from soil field capacity on oil palm still possible<sup>13</sup>, oil palm plantation with water scarcity significantly reduces growth and oil yield<sup>14,15,16</sup>.

Implication of this research is to facilitate the selection and application of fertilizer for oil palm seedlings, means that the higher the vedagro fertilizer as classified as organic fertilizer, the higher growth of oil palm seedlings is. As a recommendation of this study is effect vedagro fertilizer would be significant on oil palm seedlings at high doses of 15 g baby polybag-1, while the watering should be according to the suggestion of 0.2 liters of baby polybag-1 day-1 water.

Limitation of this study, has not been tried in multi-location dry areas in Indonesia and needs to increase response variables. CONCLUSION 1. Vedagro Fertilizer up to 15 g baby polybag-1 increase significantly plant height, but not significant effect to stem diameter, the number of leaves and leaf area of oil palm seedlings in the pre-nursery. 2.

Water stress decreases significantly in plant growth, namely plant height, stem diameter, number of leaves, leaf area of oil palm seedlings but ABA content increase significantly. Water stress can increase the ABA content as many 0.0022 ppm cm<sup>-2</sup> leaf area. 3. The interaction of doses of Vedagro fertilizer and water stress has no significant effect on plant height, stem diameter, number of leaves and leaf area of oil palm seedlings in the pre-nursery.

**SIGNIFICANCE STATEMENTS** This Study discover the effect of Vedagro fertilizer to produce better growth of oil palm seedlings and the effect of water stress on seedling growth. This study will help researcher, farmers, seed industry to uncover the critical areas on nursery in various places in Indonesia. Thus, a new theory is found that on water stress can be predicted ABA content formed, may be arrived at.

**ACKNOWLEDGMENT** We express our gratitude to the Board of Darma Agung University Foundation and Rector of Darma Agung University for their help both materially and morally. Thanks are also extended to Sitorus Lature, Gunania, Ruth Maria, Klintan Sinurat, Epianus Ndraha, Feri Sipahutar, Irwan Jaya, Fidelis Gea, Jery Frans, Sentrius Bu'ulolo for their technical assistance. **REFERENCES** Ferrianta, Y. (2017). Issue and Solution Palm Oil Industry in Indonesia. Agriculture Research and Technology 5:1-2. <https://juniperpublisher.com/artoaj/pdf/ARTOAJ.MS.ID.555656.pdf> Aba and K.P. Baiyeri, 2012. The Oil Palm. Department of Crop Science, University of Nigeria, Nsukka. [https://www.researchgate.net/publication/291331029\\_THE\\_OIL\\_PALM](https://www.researchgate.net/publication/291331029_THE_OIL_PALM) Gafur, M.A and E.T. Susila Putra, 2019. Effect of Drought Stress in Physiological Oil Palm Seedling (*Elaeis guineensis* Jacq.) using Calcium application. Science Alert.

Asian Journal of Biological Sciences. DOI: 10.3923/ajbs.2019.550.556 URL: <https://scialert.net/abstract/?doi=ajbs.2019.550.556> <https://indonesian.alibaba.com/g/vedagro-fertilizer.html>. Vedagro fertilizer brochure, 2010. 5. <https://www.google.com/search?client=firefox-bd&q=Standar+Operasional+Prosedur.+No+Dokumen%3ASOP+Agro-04%2F00>. Error! Hyperlink reference not valid.. Downloaded on April 2019. Sirait, B.A, and A.I. Manurung, 2019. Oil Palm Cultivation. Student Handbook.

The Agrotechnology Study Program, Faculty of Agriculture, University of Darma Agung Fairhurst, T and R. Hardler, 2003. Oil Palm: Management for Large and Sustainable Yield. International Potash Institute, Singapore. Mastur, 2016. Physiological Responses of Sugar Cane to Drought. Industrial Tobacco, Fiber and Oil Plant Bulletin 8 (2), October 2016: 98-111. <http://ejurnal.litbang.pertanian.go.id> > bultas > issue > download > pdf\_2 Ferreira, T.H.S, M.S.Tsunada, D.Bassi, P.Araujo, L.Mattiello, G.V. Guidelli, G.L.Righetto,

P.Lakshmanan and M.Menossi, 2017.

Sugarcane **Water Stress Tolerance Mechanisms and Its Implications on Developing Biotechnology** Solutions. *Front Plant Sci.* 2017; 8: 1077. Published online 2017 Jun 23. doi: 10.3389/fpls.2017.01077

Chaves, M.M., Maroco, J.P., and Pereira, J.S., 2003.

**Understanding plant responses to drought from genes to the whole plant.** *Functional Plant Biology*, 30. 239-264. www.publish.csiro.au/journals/fpb. DOI: 10.1071/FP02076.

Hanafi, Sirait, S.M., Purwanti E., Irawan, C., Rochaeni, H., Rosita, T., and Purba, J.A., 2017. Comparison of Abscisic Acid Hormone on the leaves of Palm Oil Seeds with Watered and not watered. *Journal of Pharmacognosy and Phytochemistry.*

www.phytojournal.com. E-ISSN:2278-4136P-ISSN:2349-8234

Caroline Salazar, Hernandez, C., Pino, M.T., 2015. Plant water stress; Associations between ethylene and abscisic acid response.

Chilean **Journal of Agricultural Research 75** (suppl.1) August 2015.

[https://scielo.conicyt.cl/scielo.php?script=sci\\_arttext&pid=S0718-8392015000300008](https://scielo.conicyt.cl/scielo.php?script=sci_arttext&pid=S0718-8392015000300008)

Najihah, T.S., M. H.Ibrahim, A.A. Razak, 2019. Effect of Water Stress on the Growth, Physiology and Biochemical Properties of Oil Palm Seedlings. *AIMS Agriculture and Food*, 2019, 4(4): 854-868. doi: 10.3934/agrfood.2019.4.854. Legros, S., Mialet-Serra, I., Caliman, J.P., Siregar, F.A., Clement-Vidal, A.,

and Dingkuhn, M. (2009). Phenology and Growth adjustmants of oil palm (*Elaeis guineensis* Jacq) to photoperiod and climate variability. *Annals of Botany*

104:1171-1182. doi:10.1093/aob/mcp214, available online at

www.aob.oxfordjournals.org Cao et al, 2011; Carr, 2011 in Duangpan, S., Sujitto, S and Eksomtramage, 2017.

**Genotypic variation in Proline Accumulation during Sequencing Drought and Rewatering in Response to Drought** Preconditioning-*International Journal of Agricultural Technology* 2017. Vol. 13 (6): 927-940. <http://www.ijat-aatsea.com>

Duangpan, S., Sujitto, S and Eksomtramage, T., 2017.

**Genotypic Variation in Proline Accumulation during Sequential Drought and Rewatering in Response to Drought** Preconditioning. *International Journal of Agricultural Technology* 2017. Vol.13 (6):927-940. Available online <http://www.ijat-aatsea.com> ISSN 2630-0192 (Online).

INTERNET SOURCES:

-----  
<1% - <https://publications.waset.org/abstracts/search?q=crude%20palm%20oil>

<1% -

[https://www.researchgate.net/publication/328347667\\_Plant\\_growth-promoting\\_rhizobacteria\\_used\\_in\\_South\\_Korea](https://www.researchgate.net/publication/328347667_Plant_growth-promoting_rhizobacteria_used_in_South_Korea)

<1% - <https://ergobiologiblog.wordpress.com/2011/11/23/ergonomic-2/>

<1% - [https://en.wikipedia.org/wiki/The\\_Lower\\_Galilee](https://en.wikipedia.org/wiki/The_Lower_Galilee)

<1% - <http://rjas.org/Newlook/Issues.aspx?Issued=43>

<1% - <https://scialert.net/fulltext/?doi=jbs.2012.146.153>

<1% -

<https://id.123dok.com/document/qogr935z-proceeding-international-seminar-on-accounting-for-society-sustainable-development-challenge-and-opportunity-for-professional-accountants-in-developing-countries-eprints-unpam.html>

<1% -

[https://archive.org/stream/NSSMHenryA.KissingerReport200435/NSSM%20Henry%20A.%20Kissinger%20Report%202004-35\\_djvu.txt](https://archive.org/stream/NSSMHenryA.KissingerReport200435/NSSM%20Henry%20A.%20Kissinger%20Report%202004-35_djvu.txt)

<1% - <http://www.gutenberg.org/files/42584/42584-h/42584-h.htm>

<1% - <https://quizlet.com/110625438/personality-exam-3-flash-cards/>

<1% - <http://www.fao.org/3/Y2783E/y2783e06.htm>

<1% - <https://www.sciencedirect.com/science/article/pii/S0254629916300990>

<1% - <https://www.mdpi.com/1422-0067/19/11/3643/htm>

1% - <https://jurnal.umsb.ac.id/index.php/agriculture/article/view/1148>

<1% -

[https://www.researchgate.net/publication/268428110\\_Morphological\\_physiological\\_and\\_biochemical\\_responses\\_of\\_plants\\_to\\_drought\\_stress](https://www.researchgate.net/publication/268428110_Morphological_physiological_and_biochemical_responses_of_plants_to_drought_stress)

<1% -

[https://www.researchgate.net/publication/328004379\\_Plant\\_growth\\_of\\_genotypes\\_of\\_Arabica\\_coffee\\_on\\_water\\_stress](https://www.researchgate.net/publication/328004379_Plant_growth_of_genotypes_of_Arabica_coffee_on_water_stress)

<1% - <http://doczz.net/doc/1724311/forest-and-biodiversity>

<1% -

[https://s3-ap-southeast-2.amazonaws.com/shared-drupal-s3fs/master-test/fapub\\_pdf/\\_R2.0/Campbelltown+Sustainable+Cities+DCP+2015+-+Volume+2+-+SUPERSEDED.pdf](https://s3-ap-southeast-2.amazonaws.com/shared-drupal-s3fs/master-test/fapub_pdf/_R2.0/Campbelltown+Sustainable+Cities+DCP+2015+-+Volume+2+-+SUPERSEDED.pdf)

<1% -

[https://www.researchgate.net/publication/281120909\\_Fungicides\\_phytotonic\\_action\\_on\\_the\\_development\\_of\\_soybean](https://www.researchgate.net/publication/281120909_Fungicides_phytotonic_action_on_the_development_of_soybean)

<1% -

<https://www.neliti.com/id/publications/223258/pengaruh-kekeringan-terhadap-pertumbuhan-dan-perkembangan-tanaman-nyamplung-call>

<1% - <https://www.sciencedirect.com/science/article/pii/S240584401838695X>

<1% - <http://journal.ipb.ac.id/index.php/jurnalagronomi/article/view/16187>

<1% - <https://www.sciencedirect.com/science/article/pii/S0378112719321772>

<1% -

<https://www.scribd.com/document/299676232/Nigerian-Journal-of-Soil-Science-Vol-25>

-2015-pdf

1% - <https://www.sciencedirect.com/science/article/pii/S2211601X1500036X>

<1% - <http://www.fao.org/3/XII/0459-B4.htm>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6248903/>

<1% - <https://quizlet.com/190395213/biology-isaempa-june-2014-flash-cards/>

<1% -

[https://jurnal.ugm.ac.id/index.php/index/oai?verb=ListRecords&metadataPrefix=oai\\_dc&set=jip:ART](https://jurnal.ugm.ac.id/index.php/index/oai?verb=ListRecords&metadataPrefix=oai_dc&set=jip:ART)

<1% -

[https://www.researchgate.net/publication/259644023\\_The\\_reaction\\_of\\_Amaranthus\\_hybridus\\_to\\_infection\\_by\\_Amaranthus\\_mosaic\\_virus](https://www.researchgate.net/publication/259644023_The_reaction_of_Amaranthus_hybridus_to_infection_by_Amaranthus_mosaic_virus)

<1% -

<https://cropandsoil.oregonstate.edu/sites/agscid7/files/crop-soil/finalseedreport2012.pdf>

<1% -

<https://www.specialmetals.com/assets/smc/documents/alloys/nickel-duranickel/nickel-200-201.pdf>

<1% -

[https://www.researchgate.net/publication/324112594\\_Simulation\\_of\\_Stomatal\\_Conductance\\_and\\_Water\\_Use\\_Efficiency\\_of\\_Tomato\\_Leaves\\_Exposed\\_to\\_Different\\_Irrigation\\_Regimes\\_and\\_Air\\_CO2\\_Concentrations\\_by\\_a\\_Modified\\_Ball-Berry\\_Model](https://www.researchgate.net/publication/324112594_Simulation_of_Stomatal_Conductance_and_Water_Use_Efficiency_of_Tomato_Leaves_Exposed_to_Different_Irrigation_Regimes_and_Air_CO2_Concentrations_by_a_Modified_Ball-Berry_Model)

<1% -

<https://www.agronomy.k-state.edu/documents/nutrient-management/nmrg-soil-nutrients,-sources-and-uptake.pdf>

<1% -

[https://www.researchgate.net/publication/288457331\\_Evaluation\\_of\\_nitrogen\\_uptake\\_efficiency\\_of\\_different\\_oil\\_palm\\_genotypes\\_using\\_15n\\_isotope\\_labelling\\_method](https://www.researchgate.net/publication/288457331_Evaluation_of_nitrogen_uptake_efficiency_of_different_oil_palm_genotypes_using_15n_isotope_labelling_method)

<1% - <http://www.nature.com/articles/srep28011>

<1% - <https://www.researchgate.net/topic/Water-Stress>

<1% - <https://bmcplantbiol.biomedcentral.com/articles/10.1186/s12870-020-02349-9>

1% -

[https://mafiadoc.com/abscisic-acid-signaling-and-abiotic-stress-tolerance-in-plants-frontiers\\_5b6bfbc8097c47d1118b45a8.html](https://mafiadoc.com/abscisic-acid-signaling-and-abiotic-stress-tolerance-in-plants-frontiers_5b6bfbc8097c47d1118b45a8.html)

<1% -

<https://repository.ipb.ac.id/bitstream/handle/123456789/62169/ACSAC-ISBN-76.pdf>

<1% - <http://docsdrive.com/pdfs/ansinet/ja/0000/70718-70718.pdf>

<1% - <https://link.springer.com/article/10.1186%2Fs11671-016-1607-2>

<1% - <https://iopscience.iop.org/issue/1755-1315/383/1>

1% -

<https://pdfs.semanticscholar.org/3e46/042424a29da1d50f1ca8e6f9fa67671d6a08.pdf>

<1% -

[https://www.researchgate.net/publication/227550299\\_Palm\\_oil\\_production\\_through\\_sustainable\\_plantations](https://www.researchgate.net/publication/227550299_Palm_oil_production_through_sustainable_plantations)

<1% - <https://id.scribd.com/doc/302609923/Laporan-Tahunan-2014>

<1% -

[https://www.researchgate.net/publication/309033558\\_Studies\\_of\\_phenotypic\\_and\\_genotypic\\_variation\\_in\\_sixteen\\_cucumber\\_genotypes](https://www.researchgate.net/publication/309033558_Studies_of_phenotypic_and_genotypic_variation_in_sixteen_cucumber_genotypes)

1% - <https://www.ncbi.nlm.nih.gov/pubmed/28690620>

<1% - <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5491614/>

<1% - <https://content.sciendo.com/view/journals/sab/50/4/article-p211.xml>

<1% - <https://scielo.conicyt.cl/pdf/chiljar/v75s1/at03.pdf>

<1% - <https://www.aimspress.com/article/10.3934/GF.2019.1.30/Related.html>

<1% -

[https://www.researchgate.net/publication/231922452\\_THE\\_water\\_relations\\_and\\_irrigation\\_requirements\\_of\\_oil\\_palm\\_Elaeis\\_guineensis\\_A\\_review](https://www.researchgate.net/publication/231922452_THE_water_relations_and_irrigation_requirements_of_oil_palm_Elaeis_guineensis_A_review)

1% - <https://grad.psu.ac.th/th/faculty-staff/lecturer.html?tid=2772>

<1% - <http://ijat-aatsea.com/overview.html>